

1. Baird L, Yamamoto M. The Molecular Mechanisms Regulating the KEAP1–NRF2 Pathway. *Mol Cell Biol.* 2020;40(13). doi:10.1128/MCB.00099–20
2. Watahiki A, Shimizu K, Hoshikawa S, et al. Lipin–2 degradation elicits a proinflammatory gene signature in macrophages. *Biochem Biophys Res Commun.* 2020;524(2):477–483. doi:10.1016/j.bbrc.2020.01.119
3. Vos RA, Katayama T, Mishima H, et al. BioHackathon 2015: Semantics of data for life sciences and reproducible research. *F1000Research.* 2020;9:136. doi:10.12688/f1000research.18236.1
4. Yaoita N, Satoh K, Satoh T, et al. Identification of the Novel Variants in Patients With Chronic Thromboembolic Pulmonary Hypertension. *J Am Heart Assoc.* 2020;9(21). doi:10.1161/JAHA.120.015902
5. Nishizawa A, Kumada K, Tateno K, et al. Analysis of HLA–G long–read genomic sequences in mother–offspring pairs with preeclampsia. *Sci Rep.* 2020;10(1):20027. doi:10.1038/s41598–020–77081–3
6. Saigusa D, Motoike IN, Saito S, et al. Impacts of NRF2 activation in non–small–cell lung cancer cell lines on extracellular metabolites. *Cancer Sci.* 2020;111(2):667–678. doi:10.1111/cas.14278
7. Baird L, Suzuki T, Takahashi Y, Hishinuma E, Saigusa D, Yamamoto M. Geldanamycin–Derived HSP90 Inhibitors Are Synthetic Lethal with NRF2. *Mol Cell Biol.* 2020;40(22). doi:10.1128/MCB.00377–20
8. Tsuboi A, Matsui H, Shiraishi N, et al. Design and Progress of Oral Health Examinations in the Tohoku Medical Megabank Project. *Tohoku J Exp Med.* 2020;251(2):97–115. doi:10.1620/tjem.251.97
9. Lin Y, Nakatochi M, Hosono Y, et al. Genome–wide association meta–analysis identifies GP2 gene risk variants for pancreatic cancer. *Nat Commun.* 2020;11(1):3175. doi:10.1038/s41467–020–16711–w
10. Takahashi Y, Ueki M, Yamada M, et al. Improved metabolomic data–based prediction of depressive symptoms using nonlinear machine learning with feature selection. *Transl Psychiatry.* 2020;10(1):157. doi:10.1038/s41398–020–0831–9
11. Matsunaga H, Ito K, Akiyama M, et al. Transethnic Meta–Analysis of Genome–Wide Association Studies Identifies Three New Loci and Characterizes Population–Specific Differences for Coronary Artery Disease. *Circ Genomic Precis Med.* 2020;13(3). doi:10.1161/CIRCGEN.119.002670
12. Kuriyama S, Metoki H, Kikuya M, et al. Cohort Profile: Tohoku Medical Megabank Project Birth and Three–Generation Cohort Study (TMM BirThree Cohort Study): rationale, progress and perspective. *Int J Epidemiol.* 2020;49(1):18–19m. doi:10.1093/ije/dyz169

13. Oishi T, Matsumaru D, Ota N, et al. Activation of the NRF2 pathway in Keap1-knockdown mice attenuates progression of age-related hearing loss. *npj Aging Mech Dis.* 2020;6(1):14. doi:10.1038/s41514-020-00053-4
14. Okazaki K, Anzawa H, Liu Z, et al. Enhancer remodeling promotes tumor-initiating activity in NRF2-activated non-small cell lung cancers. *Nat Commun.* 2020;11(1):5911. doi:10.1038/s41467-020-19593-0
15. Wada Y, Kikuchi A, Arai-Ichinoi N, et al. Correction: Biallelic GALM pathogenic variants cause a novel type of galactosemia. *Genet Med.* 2020;22(7):1281. doi:10.1038/s41436-020-0836-z
16. Akiyama T, Saigusa D, Hyodo Y, et al. Metabolic Profiling of the Cerebrospinal Fluid in Pediatric Epilepsy. *Acta Med Okayama.* 2020;74(1):65–72. doi:10.18926/AMO/57955
17. Kasai T, Ono S, Koshiba S, et al. Amino-acid selective isotope labeling enables simultaneous overlapping signal decomposition and information extraction from NMR spectra. *J Biomol NMR.* 2020;74(2–3):125–137. doi:10.1007/s10858-019-00295-9
18. Wagata M, Ishikuro M, Obara T, et al. Low birth weight and abnormal pre-pregnancy body mass index were at higher risk for hypertensive disorders of pregnancy. *Pregnancy Hypertens.* 2020;22:119–125. doi:10.1016/j.preghy.2020.08.001
19. Narita A, Nagai M, Mizuno S, et al. Clustering by phenotype and genome-wide association study in autism. *Transl Psychiatry.* 2020;10(1):290. doi:10.1038/s41398-020-00951-x
20. Takahashi Y, Ueki M, Tamiya G, et al. Machine learning for effectively avoiding overfitting is a crucial strategy for the genetic prediction of polygenic psychiatric phenotypes. *Transl Psychiatry.* 2020;10(1):294. doi:10.1038/s41398-020-00957-5
21. Gutiérrez Rico EM, Kikuchi A, Saito T, et al. CYP2D6 genotyping analysis and functional characterization of novel allelic variants in a Ni-Vanuatu and Kenyan population by assessing dextromethorphan O-demethylation activity. *Drug Metab Pharmacokinet.* 2020;35(1):89–101. doi:10.1016/j.dmpk.2019.07.003
22. Koshiba S, Motoike IN, Saigusa D, et al. Identification of critical genetic variants associated with metabolic phenotypes of the Japanese population. *Commun Biol.* 2020;3(1):662. doi:10.1038/s42003-020-01383-5
23. Hishinuma E, Gutiérrez Rico E, Hiratsuka M. In Vitro Assessment of Fluoropyrimidine-Metabolizing Enzymes: Dihydropyrimidine Dehydrogenase, Dihydropyrimidinase, and β -Ureidopropionase. *J Clin Med.* 2020;9(8):2342. doi:10.3390/jcm9082342
24. Taguchi K, Yamamoto M. The KEAP1-NRF2 System as a Molecular Target of Cancer Treatment. *Cancers (Basel).* 2020;13(1):46. doi:10.3390/cancers13010046

25. Nagai K, Uranbileg B, Chen Z, et al. Identification of novel biomarkers of hepatocellular carcinoma by high-definition mass spectrometry: Ultrahigh-performance liquid chromatography quadrupole time-of-flight mass spectrometry and desorption electrospray ionization mass spectrometry imaging. *Rapid Commun Mass Spectrom.* 2020;34(S1):e8551. doi:10.1002/rcm.8551
26. Kojima K, Tadaka S, Katsuoka F, Tamiya G, Yamamoto M, Kinoshita K. A genotype imputation method for de-identified haplotype reference information by using recurrent neural network. Ay F, ed. *PLOS Comput Biol.* 2020;16(10):e1008207. doi:10.1371/journal.pcbi.1008207
27. Suzuki T, Uruno A, Yumoto A, et al. Author Correction: Nrf2 contributes to the weight gain of mice during space travel. *Commun Biol.* 2020;3(1):566. doi:10.1038/s42003-020-01292-7
28. Suzuki T, Uruno A, Yumoto A, et al. Nrf2 contributes to the weight gain of mice during space travel. *Commun Biol.* 2020;3(1):496. doi:10.1038/s42003-020-01227-2
29. Ninomiya K, Takatsuki T, Kushida T, Yamamoto Y, Ogishima S. Choosing preferable labels for the Japanese translation of the Human Phenotype Ontology. *Genomics Inform.* 2020;18(2):e23. doi:10.5808/GI.2020.18.2.e23
30. Yokose T, Kabe Y, Matsuda A, et al. O-Glycan-Altered Extracellular Vesicles: A Specific Serum Marker Elevated in Pancreatic Cancer. *Cancers (Basel).* 2020;12(9):2469. doi:10.3390/cancers12092469
31. Ishida N, Aoki Y, Katsuoka F, et al. Landscape of electrophilic and inflammatory stress-mediated gene regulation in human lymphoblastoid cell lines. *Free Radic Biol Med.* 2020;161:71–83. doi:10.1016/j.freeradbiomed.2020.09.023
32. Hayashi M, Kuga A, Suzuki M, et al. Microenvironmental Activation of Nrf2 Restricts the Progression of Nrf2-Activated Malignant Tumors. *Cancer Res.* 2020;80(16):3331–3344. doi:10.1158/0008-5472.CAN-19-2888
33. Kumondai M, Hishinuma E, Gutiérrez Rico EM, et al. Heterologous expression of high-activity cytochrome P450 in mammalian cells. *Sci Rep.* 2020;10(1):14193. doi:10.1038/s41598-020-71035-5
34. Katayama T, Kawashima S, Micklem G, et al. BioHackathon series in 2013 and 2014: improvements of semantic interoperability in life science data and services. *F1000Research.* 2019;8:1677. doi:10.12688/f1000research.18238.1
35. Saito T, Kuma A, Sugiura Y, et al. Autophagy regulates lipid metabolism through selective turnover of NCoR1. *Nat Commun.* 2019;10(1):1567. doi:10.1038/s41467-019-08829-3
36. Shimizu M, Yoda H, Nakakuki K, et al. Genetic variants of flavin-containing monooxygenase 3 (FMO3) derived from Japanese subjects with the trimethylaminuria phenotype and whole-genome sequence data from a large Japanese database. *Drug Metab Pharmacokinet.* 2019;34(5):334–339. doi:10.1016/j.dmpk.2019.06.001

37. Larsson SC, Traylor M, Burgess S, et al. Serum magnesium and calcium levels in relation to ischemic stroke. *Neurology*. 2019;92(9). doi:10.1212/WNL.0000000000007001
38. Dodo M, Kitamura H, Shima H, et al. Lactate dehydrogenase C is required for the protein expression of a sperm-specific isoform of lactate dehydrogenase A. *J Biochem*. 2019;165(4):323–334. doi:10.1093/jb/mvy108
39. Tanikawa C, Kamatani Y, Terao C, et al. Novel Risk Loci Identified in a Genome-Wide Association Study of Urolithiasis in a Japanese Population. *J Am Soc Nephrol*. 2019;30(5):855–864. doi:10.1681/ASN.2018090942
40. Sugawara J, Ochi D, Yamashita R, et al. Maternity Log study: a longitudinal lifelog monitoring and multiomics analysis for the early prediction of complicated pregnancy. *BMJ Open*. 2019;9(2):e025939. doi:10.1136/bmjopen-2018-025939
41. Malik R, Chauhan G, Traylor M, et al. Publisher Correction: Multiancestry genome-wide association study of 520,000 subjects identifies 32 loci associated with stroke and stroke subtypes. *Nat Genet*. 2019;51(7):1192–1193. doi:10.1038/s41588-019-0449-0
42. Yasuda J, Kinoshita K, Katsuoka F, et al. Genome analyses for the Tohoku Medical Megabank Project towards establishment of personalized healthcare. *J Biochem*. 2019;165(2):139–158. doi:10.1093/jb/mvy096
43. Wada Y, Kikuchi A, Arai-Ichinoi N, et al. Biallelic GALM pathogenic variants cause a novel type of galactosemia. *Genet Med*. 2019;21(6):1286–1294. doi:10.1038/s41436-018-0340-x
44. Sato T, Kawasaki Y, Maekawa M, et al. Value of global metabolomics in association with diagnosis and clinicopathological factors of renal cell carcinoma. *Int J Cancer*. 2019;145(2):484–493. doi:10.1002/ijc.32115
45. Yamaguchi-Kabata Y, Yasuda J, Uruno A, et al. Estimating carrier frequencies of newborn screening disorders using a whole-genome reference panel of 3552 Japanese individuals. *Hum Genet*. 2019;138(4):389–409. doi:10.1007/s00439-019-01998-7
46. Minegishi N, Nishijima I, Nobukuni T, et al. Biobank Establishment and Sample Management in the Tohoku Medical Megabank Project. *Tohoku J Exp Med*. 2019;248(1):45–55. doi:10.1620/tjem.248.45
47. Nagasaki M, Kuroki Y, Shibata TF, et al. Construction of JRG (Japanese reference genome) with single-molecule real-time sequencing. *Hum Genome Var*. 2019;6(1):27. doi:10.1038/s41439-019-0057-7
48. Shido K, Kojima K, Yamasaki K, et al. Susceptibility Loci for Tanning Ability in the Japanese Population Identified by a Genome-Wide Association Study from the Tohoku Medical Megabank Project Cohort Study. *J Invest Dermatol*. 2019;139(7):1605–1608.e13. doi:10.1016/j.jid.2019.01.015

49. Sakurai-Yageta M, Kawame H, Kuriyama S, et al. A training and education program for genome medical research coordinators in the genome cohort study of the Tohoku Medical Megabank Organization. *BMC Med Educ.* 2019;19(1):297. doi:10.1186/s12909-019-1725-5
50. Mizuguchi T, Suzuki T, Abe C, et al. A 12-kb structural variation in progressive myoclonic epilepsy was newly identified by long-read whole-genome sequencing. *J Hum Genet.* 2019;64(5):359-368. doi:10.1038/s10038-019-0569-5
51. Tadaka S, Katsuoka F, Ueki M, et al. 3.5KJPNv2: an allele frequency panel of 3552 Japanese individuals including the X chromosome. *Hum Genome Var.* 2019;6(1):28. doi:10.1038/s41439-019-0059-5
52. Watanabe T, Saito T, Rico EMG, et al. Functional characterization of 40 CYP2B6 allelic variants by assessing efavirenz 8-hydroxylation. *Biochem Pharmacol.* 2018;156:420-430. doi:10.1016/j.bcp.2018.09.010
53. Saito T, Gutiérrez Rico EM, Kikuchi A, et al. Functional characterization of 50 CYP2D6 allelic variants by assessing primaquine 5-hydroxylation. *Drug Metab Pharmacokinet.* 2018;33(6):250-257. doi:10.1016/j.dmpk.2018.08.004
54. Kumondai M, Ito A, Hishinuma E, et al. Development and application of a rapid and sensitive genotyping method for pharmacogene variants using the single-stranded tag hybridization chromatographic printed-array strip (STH-PAS). *Drug Metab Pharmacokinet.* 2018;33(6):258-263. doi:10.1016/j.dmpk.2018.08.003